

# BIBLIOGRAPHY

## CULTURAL COMPETENCY RESOURCES

---

### Assessment Tools

Campinha-Bacote J (1998). Inventory for Assessing the Process of Cultural Competence (IAPCC) Among Health Care Professionals. Cincinnati, OH: Transcultural C.A.R.E. Associates, 11108 Huntwicke Place, Cincinnati 45241, 513-469-1664.

Dana RH, Behn JD, Gonwa T (1992). A Checklist for the Examination of Cultural Competence in Social Service Agencies in Research on Social Work Practice, Vol. 2, No.2, 220-233.

Mason JL (1995). The Cultural Competence Self-assessment Questionnaire: A Manual for Users. Portland Research and Training Center, Portland, Oregon.  
National Public Health and Hospital Institute (1997).

Self-Assessment of Cultural Competence. Washington, DC, NPHHI, 1212 New York Ave NW, Ste 800, Washington, DC 20005, phone 202-408-0229, fax 202-408-0235.

Success by 6/United Way of Minneapolis and Hennepin Medical Society (1996). Cultural Competence Clinic Assessment Tool. Minneapolis, MN. Center for Cross-Cultural Health, W-227, 410 Church Street, Minneapolis, MN 55455.

Taylor, Tawara D. Cultural Competence in Primary Health Care: Self-Assessment. Georgetown University Child Development Center, 3800 Reservoir Road, NW, Washington, DC 20007, 202-687-8635.

Tirado, Miguel D. (1996). Tools for Monitoring Cultural Competence in Health Care. Latino Coalition for a Healthy California, San Francisco, CA. 1535 Mission Street, San Francisco, CA 94103, 415-431-7430.

Tripp-Reimer T, Brink PJ, Saunders JM (1984). Cultural Assessment: Content and Process. Nursing Outlook, Vol. 32, No. 2, 78-82.

### Training Manuals

Cultural Competency in Health Care: A Guide for Trainers (Second Edition), 1999, National Multicultural Institute Publications, 202-483-0700 or <http://www.nmci.org>

Cultural Friendly TB Care, Developing Skills to Serve Diverse Ethnic Communities: An Asian Pacific Islander Example, October 1998. Coalition Against Tuberculosis Los Angeles, c/o American Lung Association of Los Angeles County, 323-935-5864.

## CONDUCTING NEEDS ASSESSMENT AND EVALUATION

---

Ashbaugh JW, Hoff MK, Schneider LC, Goldstrom ID, Manderscheid RW. Conducting state and local health need assessments. Designs and strategies. *Evaluation and the Health Professions*. 1983;6(4):413-434.

Devine AE. Community assessment: its components and significance in planning health care programs. *Journal of Enterostomal Therapy*. 1985;12(5):167-174.

Dignan, MB, Carr, PA. *Program Planning for Health Education and Promotion*. 2<sup>nd</sup> ed. Philadelphia, PA: Lea & Febiger;1992.

Folch-Lyon E, Trost JF. Conducting Focus Group Sessions. *Studies in Family Planning*. 1981;12(12):443-449.

Making Health Communication Programs Work: A Planner's Guide. US Department of Health and Human Services, Public Health Services, National Institutes of Health. *NIH Publication No. 92-1493*. April 1992.

Needs Assessment for HIV/AIDS Prevention and Service Programs. United States Conference of Mayors. *Technical Assistance Reports, USCM HIV Program*. Washington, DC. September 1993.

Quinn DC. Proving the effectiveness of community programs in jeopardy: the role of formative evaluation. *Quality Management in Health Care*. 2001;9(4):33-46.

Trocchio J. The hows and whys of conducting a community needs assessment. *Trustee*. 1994;47(3):6-7,27.

Valdiserri RO, West GR. Barriers to the assessment of unmet need in planning HIV/AIDS prevention programs. *Public Administration Review*. 1994;54(1):25-30.

## DIRECTLY OBSERVED THERAPY

---

Bayer R, Stayton S, Desvarieux M, Heaton C, Landesman S, Tsai WY. Directly observed therapy and treatment completion for tuberculosis in the United States: Is universal supervised therapy necessary? *American Journal of Public Health*. 1998;88(7):1052-1058.

Centers for Disease Control and Prevention. Approaches to improving adherence to antituberculosis therapy – South Carolina and New York, 1986-1991. *Morbidity and Mortality Weekly Report*. 1993;42(04):74-75,81.

Chaisson RE, Barnes GL, Hackman J, Watkinson L, Kimbrough L, Metha S, Cavalcante S, Moore RD. A randomized, controlled trial of interventions to improve adherence to isoniazid therapy to prevent tuberculosis in injection drug users. *American Journal of Medicine*. 2001;110(8):610-615.

Chaulk CP, Kazandjian VA. For the Public Health Tuberculosis Guidelines Panel. Directly observed therapy for treatment completion of pulmonary tuberculosis. *Journal of the American Medical Association*. 1998;279:943-948.

Gourevitch MN, Alcabes P, Wasserman WC, Arno PS. Cost-effectiveness of directly observed chemoprophylaxis of tuberculosis among drug users at high risk for tuberculosis. *International Journal of Tuberculosis and Lung Disease*. 1998;2(7):531-540.

Jasmer RM. Treatment of latent tuberculosis infection. *Tuberculosis Case Management and Contact Investigation 2001 Course Binder*. Francis J. Curry National Tuberculosis Center, San Francisco, CA. 2001.

## **HARM REDUCTION RESOURCES**

---

Bigg D. Substance use management: a harm reduction-principled approach to assisting the relief of drug-related problems. *Journal of Psychoactive Drugs*. 2001;33(1):33-38.

Denning P. Strategies for implementation of harm reduction in treatment settings. *Journal of Psychoactive Drugs*. 2001;33(1):23-26.

Denning P, Little J. Harm reduction in mental health – The emerging work of harm reduction psychotherapy. *Harm Reduction Communication*. Spring 2001; Issue No. 11:7-10.

Fisk, SN. The association of nurses in AIDS care. Position paper on harm reduction and HIV care for drug users: integrating harm-reduction methods and HIV care. *Journal of the Association of Nurses in AIDS Care*. 1998;9(3):19-24.

Gleghorn A, Rosenbaum M, Garcia BA. Bridging the gap in San Francisco: The process of integrating harm reduction and traditional substance abuse services. *Journal of Psychoactive Drugs*. 2001;33(1):1-7.

Gunn N, White C, Srinivasan R. Primary care as harm reduction for injection drug users. *JAMA*. 1998;280:1191-1195.

Marlatt GA. Harm reduction: come as you are. *Addictive Behaviors*. 1996;21(6):779-788.

National Association for Public Health Policy. A public health approach to mitigating the negative consequences of illicit drug use. *Journal of Public Health Policy*. 1999;20(3):268-281.

## HIV AND LATENT TUBERCULOSIS INFECTION

---

Gordin et al. A controlled trial of isoniazid in persons with anergy and human immunodeficiency virus infection who are at high risk for tuberculosis. *The New England Journal of Medicine*. 1997;337(5):315.

Moreno, et al. Isoniazid preventive therapy in human immunodeficiency virus-infected persons. *Archives of Internal Medicine*. 1997;157(15):1729.

## USE OF INCENTIVES AND ENABLERS

---

Centers for Disease Control and Prevention. Approaches to improving adherence to antituberculosis therapy – South Carolina and New York, 1986-1991. *Morbidity and Mortality Weekly Report*. 1993;42(04):74-75,81.

Eastman SJ, Judd K. The creative use of incentives in TB control and prevention. *Training Files*. Francis J. Curry National Tuberculosis Center, San Francisco, CA. 1994.

Goldberg SV, Friedly C, Nolan CM. Program Operations and Evaluation Incentives in TB Treatment. *Public Health, Seattle-King County*. Seattle, WA.

Malotte CK, Hollingshead JR, Larro M. Incentives vs. outreach workers for latent tuberculosis treatment in drug users. *American Journal of Preventive Medicine*. 2001;20(2):103-107.

Moore RD, Chaulk CP, Griffiths R, Cavalcante S, Chaisson RE. Cost-effectiveness of directly observed versus self-administered therapy for tuberculosis. *American Journal of Respiratory and Critical Care Medicine*. 1996;154(4):1013-1019.

Snyder DC, Chin DP. Cost-effectiveness analysis of directly observed therapy for patients with tuberculosis at low risk for treatment default. *American Journal of Respiratory and Critical Care Medicine*. 1999;160(2):582-586.

## ADDITIONAL BACKGROUND RESOURCES

---

Adapting Guidelines for Treatment of LTBI for Local Jurisdictions (session handouts). California Tuberculosis Controllers Association. Spring 2001.

Buskin SE, Gale JL, Weiss NS, Nolan CM. Tuberculosis risk factors in adults in King County, Washington, 1988 through 1990. *American Journal of Public Health*. 1994;84(11):1750-1756.

Centers for Disease Control and Prevention. Essential components of a tuberculosis prevention and control program. Recommendations of the Advisory Council for the elimination of tuberculosis. *Morbidity and Mortality Weekly Report*. 1995;44(RR-11):1-16.

Centers for Disease Control and Prevention. Prevention and control of tuberculosis among homeless persons. Recommendations of the Advisory Council for the elimination of tuberculosis. *Morbidity and Mortality Weekly Report*. 1992;41(RR-5).

Centers for Disease Control and Prevention. Prevention and control of tuberculosis in U.S. communities with at-risk minority populations. Recommendations of the Advisory Council for the elimination of tuberculosis. *Morbidity and Mortality Weekly Report*. 1992;41(RR-5):1-12.

Centers for Disease Control and Prevention. Prevention and treatment of tuberculosis among patients infected with human immunodeficiency virus: principles of therapy and revised recommendations. *Morbidity and Mortality Weekly Report*. 1998;47(RR-20):1-58.

Centers for Disease Control and Prevention. Targeted tuberculin testing and treatment of latent tuberculosis infection. *Morbidity and Mortality Weekly Report*. 2000;49(RR-6):1-51.

Friedman LN, Williams MT, Singh TP, Frieden TR. Tuberculosis, AIDS, and death among substance abusers on welfare in New York City. *The New England Journal of Medicine*. 1996;334(13):828-833.

Gaiter L, ed. Ending neglect: the elimination of tuberculosis in the United States. Institute of Medicine. National Academy Press, Washington, D.C. 2000.

Gordin et al. A controlled trial of isoniazid in persons with anergy and human immunodeficiency virus infection who are at high risk for tuberculosis. *The New England Journal of Medicine*. 1997;337(5):315.

Gourevitch MN, Hartel D, Selwyn PA, Schoenbaum EE, Klein RS. Effectiveness of isoniazid chemoprophylaxis for HIV-infected drug users at high risk for active tuberculosis. *AIDS*. 1999;13(15):2069-2074.

Malotte CK, Rhodes F, Mais KE. Tuberculosis screening and compliance with return for skin test reading among active drug users. *American Journal of Public Health*. 1998;88(5):792-796.

Markowitz N, Hansen NI, Hopewell PC, Glassroth J, Kvale PA, Mangura BT, Wilcosky TC, Wallace JM, Rosen MJ, Reichman LB, and the Pulmonary Complications of HIV Infection Study Group. Incidence of tuberculosis in the United States among HIV-infected persons. *Annals of Internal Medicine*. 1997;126:123-132.

Moreno et al. Isoniazid preventive therapy in human immunodeficiency virus-infected persons. *Archives of Internal Medicine*. 1997;157(15):1729.

Morse G. A review of case management for people who are homeless: implications for practice, policy, and research. Office of the Assistant Secretary for Planning and Evaluation, U.S. Department of Health and Human Services. Downloaded from - <http://aspe.hhs.gov/search/progsys/homeless/symposium/7-Casemgmt.htm>

National TB program objectives. Program Announcement 00001 – Tuberculosis elimination and laboratory cooperative agreement (also referred to as the “FY 2000 TB Cooperative Agreement”).

Rosenheck R, Bassuk E, Salomon A. Special populations of homeless Americans. Office of the Assistant Secretary for Planning and Evaluation, U.S. Department of Health and Human Services. Downloaded from - <http://aspe.hhs.gov/search/progsys/homeless/symposium/2-Spclpop.htm>

Working with patients with special challenges. DOT Trainer’s Curriculum, Session #5. Francis J. Curry National Tuberculosis Center, San Francisco. 2001.